

CCD Photometry of a Newly Confirmed SU UMa-Type Dwarf Nova, NSV 4838

Akira IMADA, Tatsuki YASUDA, Toshihiro OMODAKA, Shota OIZUMI, Hiroyuki YAMAMOTO, Shunsuke TANADA,
 Yoshihiro ARAO, Kie KODAMA, Miho SUZUKI, Takeshi MATSUO

Faculty of Science, Kagoshima University, 1-21-30 Korimoto, Kagoshima-shi, Kagoshima 890-0065
yasuda@astro.sci.kagoshima-u.ac.jp

Hiroyuki MAEHARA, Taichi KATO, Kei SUGIYASU, Yuuki MORITANI, Masanao SUMIYOSHI
Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502

Kazuhiro NAKAJIMA
VSOLJ, 124 Isatotyo Teradani, Kumano, Mie 519-4673
 John PIETZ

Rostocker Str. 62, 50374 Erfstadt, Germany
 Kenshi YANAGISAWA

Okayama Astrophysical Observatory, National Astronomical Observatory of Japan, Asakuchi, Okayama 719-0232
 and

Daisaku NOGAMI
Hida Observatory, Kyoto University, Kamitakara, Gifu 506-1314

(Received (reception date); accepted (acceptation date))

Abstract

We present time-resolved CCD photometry of a dwarf nova NSV 4838 (UMa 8, SDSS J102320.27+440509.8) during the 2005 June and 2007 February outburst. Both light curves showed superhumps with a mean period of 0.0699(1) days for the 2005 outburst and 0.069824(83) days for the 2007 outburst, respectively. Using its known orbital period of 0.0678 days, we estimated the mass ratio of the system to be $q=0.13$ based on an empirical relation. Although the majority of SU UMa-type dwarf novae having similar superhump periods show negative period derivatives, we found that the superhump period increased at $\dot{P}/P_{\text{sh}}=+7(+3, -4)\times 10^{-5}$ during the 2007 superoutburst. We also investigated long-term light curves of NSV 4838, from which we derived 340 days as a supercycle of this system.

Key words: accretion, accretion disks — stars: dwarf novae — stars: individual(NSV 4838) — stars: novae, cataclysmic variables

1. Introduction

Dwarf novae are a group of cataclysmic variable stars that consist of a white dwarf (primary) and a late-type star (secondary) (for a review, see e.g., Warner 1995; Osaki 1996; Hellier 2001; Osaki 2005; Connon Smith 2007). The orbital periods of the majority of dwarf novae are below 9 hours (Gänsicke 2005). The secondary star fills its Roche lobe, transferring gas into the primary Roche lobe via the inner Lagrangian point (L1). This process results in formation of an accretion disk around the primary. The accretion disk plays the main role in dwarf nova outbursts which are driven by the thermal limit-cycle instability (Hoshi 1979; Meinunger 1981; Cannizzo et al. 1988).

SU UMa-type dwarf novae are one subclass of dwarf novae, whose orbital periods are shorter than 2 hours except for a few systems (for a review, see e.g., Osaki 1989). These systems exhibit two types of outbursts. One is normal outburst, whose duration is typically a few days. The other is superoutburst, whose duration is longer than 10 days and their maximum magnitude is ~ 0.5 mag brighter than that of normal outburst. During the superoutburst,

the light curve shows tooth-like modulations called superhump with an amplitude of ~ 0.2 mag. The period of the superhump (P_{sh}) is a few percent longer than those of the orbital period (P_{orb}). This slightly longer periodicity and the modulations are attributed to phase-dependent removal of the angular momentum in a tidally deformed eccentric disk (Whitehurst 1988; Hirose, Osaki 1990; Murray 1998; Smith et al. 2007).

NSV 4838 (UMa 8 (Downes et al. 2001), SDSS J102320.27+440509.8 (Szkody et al. 2005)) was listed by Berger, Fringant (1977) as a class I (UV-richest) object. Downes et al. (1997) introduced this object as a candidate dwarf nova or a nova-like star with its magnitude ranging 14.5p-16.5p. An optical spectrum was published by Szkody et al. (2005), in which Balmer emission lines are superimposed on a blue continuum. Szkody et al. (2005) also gave the SDSS quiescent magnitudes of the object as $u=18.513$, $g=18.830$, $r=18.688$, $i=18.570$, and $z=18.514$. Radial velocity studies were implemented by Thorstensen et al. (unpublished), who determined $P_{\text{orb}}=0.0678$ days (see also, Ritter, Kolb 2003). No 2MASS counterpart is detected, but recent infrared photometry using the OAO/ISLE (Yanagisawa et al. 2006)

Table 2. List of observers.

ID	Observer	Site	Telescope
JP	J. Pietz	Erftstadt, Germany	20cm
KU	T. Yasuda+*	Kagoshima, Japan	100cm
Kyoto	A. Imada+†	Kyoto, Japan	40cm
Mhh	H. Maehara	Saitama, Japan	25cm
Njh	K. Nakajima	Mie, Japan	25cm

*T. Yasuda, S. Oizumi, H. Yamamoto, S. Tanada, Y. Arao, K. Kodama, M. Suzuki, and T. Matsuo.

†A. Imada, T. Kato, and K. Sugiyasu.

yields $H=17.51(0.09)$ and $K_s=16.81(0.08)$, respectively. The SU UMa nature of the object was initially confirmed by T. Vanmunster by detection of superhumps during the 2005 June superoutburst. Here we report on photometric studies of the 2005 June and 2007 February superoutbursts, as well as the long-term behavior of the object.

2. Observations

Time-resolved CCD photometry was performed from 2005 June 6 to June 14, and from 2007 February 7 to 28 at 5 sites. The log of observations is summarized in table 1. Detailed information on sites is given in table 2. The total data points amounted to 7428. The exposure times were 30-180 seconds without filter. All CCD systems are close to the Kron-Cousins R_C band. After debiasing and flat-fielding, we performed aperture photometry using IRAF¹ apphot for data of KU and Mhh, AIP4WIN (Berry, Burnell 2000) for data of JP, and FitsPhot4.1² for data of Njh, respectively. The Kyoto team (Kyoto) used a Java-based point spread function (PSF) photometry package developed by one of the authors (TK). The differential magnitude among each site was adjusted to that of the KU site, where we chose the comparison star NOMAD 1340-0218270, RA:10:23:33.34, Dec:+44:03:26.0 $V=15.20$, $R=15.16$). The constancy of the star is checked by nearby stars in the same frame. A heliocentric correction was made for each dataset before the following analyses.

3. Results

3.1. 2007 February superoutburst

3.1.1. light curve

The light curve obtained during the 2007 February superoutburst is shown in Figure 1. The plateau phase lasted at least 12 days³ and the variable almost linearly faded at a rate of $0.10(1) \text{ mag d}^{-1}$. After the plateau




Fig. 1. Averaged light curve of the 2007 February superoutburst. The horizontal and vertical axes indicate the HJD and the relative magnitude, respectively. The filled circles represent the averaged magnitudes for each run. The size of the standard error bar is comparable to that of the datapoints. The magnitude obtained on HJD 2454141 deviates slightly from the general trend, possibly due to sparse data and an unfavorable weather. A hint of a rebrightening was seen on HJD 2454157. The magnitude of the comparison star is $V=15.20$.

stage, NSV 4838 entered the rapid decline phase at a rate of $0.86(1) \text{ mag d}^{-1}$. These values were typical for the superoutburst of SU UMa-type dwarf novae. Although the data were noisy during the post outburst stage, a hint of a rebrightening was seen on HJD 2454157.

3.1.2. superhump

In order to estimate the mean superhump period during the plateau stage, we performed the phase dispersion minimization method (PDM, Stellingwerf 1978) using 6750 points between 2007 February 7 and 19. Figure 2 shows the resultant theta diagram, indicating that the best estimated period is $0.069824(83)$ days. The error of the period was estimated using the Lafler-Kinman class of methods (Ferne 1989). Daily averaged light curves folded by this period are displayed in figure 3. Also shown is a representative light curve on 2007 February 12 (HJD 2454143.9454-2454144.3882) obtained at KU (figure 4). No evidence for an eclipse indicates a low-to-mid inclination of NSV 4838.

3.1.3. superhump period change

We measured maximum timings of superhumps, which is summarized in table 3. A linear regression yields the following equation on the superhump maximum timings:

$$HJD(max) = 2454139.2590(24) + 0.069802(21) \times E. \quad (1)$$

Using this equation and table 3, we obtained the $O-C$ diagram displayed in figure 5. The solid curve in this figure indicates the best fitting quadratic equation between $E=0$ and 101, which is given by

$$O-C = -3.75(1.11) \times 10^{-3} - 1.62(0.41) \times 10^{-4}E + 3.09(0.40) \times 10^{-6}E^2. \quad (2)$$

superoutburst.


¹ IRAF (Image Reduction and Analysis Facility) is distributed by US National Optical Observatories, which is operated by the Association of Universities, for Research in Astronomy, Inc., under cooperative agreement with National Science Foundation.

² http://www.geocities.jp/nagai_kazuo/dload-1.html

³ The last observation before this outburst was performed at JD 2454125.8414, when the star was fainter than 16.2 in visual magnitude. Thus we were unable to constrain the duration of the

Table 1. Log of observation.

Date	HJD(start)*	HJD(end)*	N [†]	Exp [‡]	ID [§]
2005 Jun 6	53528.3834	53528.4586	46	45-60	JP
2005 Jun 7	53529.3923	53529.5135	96	45-60	JP
2005 Jun 8	53530.3811	53530.5200	107	45-60	JP
2005 Jun 9	53531.3804	53531.5176	111	45-60	JP
2005 Jun 13	53535.3978	53535.4050	6	60	JP
2005 Jun 14	53536.4019	-	1	60	JP
2007 Feb 7	54139.2154	54139.3614	70	180	KU
	54139.2237	54139.3387	210	30	Njh
2007 Feb 9	54141.2600	54141.2859	19	30	Njh
2007 Feb 10	54142.2642	54142.3415	35	180	KU
2007 Feb 11	54142.9669	54143.1278	76	180	KU
	54143.0996	54143.3036	912	30	Mhh
	54143.1802	54143.3029	224	30	Njh
2007 Feb 12	54143.9454	54144.3882	189	180	KU
	54144.1465	54144.3098	844	30	Mhh
	54144.1493	54144.3065	288	30	Njh
2007 Feb 14	54146.1660	54146.3038	139	30	Njh
	54146.2009	54146.3616	63	180	KU
2007 Feb 15	54147.1399	54147.3053	850	30	Mhh
2007 Feb 16	54148.0700	54148.2768	445	30	Kyoto
	54148.1642	54148.3079	748	30	Mhh
2007 Feb 18	54150.0437	54150.2397	1028	30	Mhh
	54150.2121	54150.3842	79	180	KU
2007 Feb 19	54150.9889	54151.1981	457	30	Kyoto
	54151.2283	54151.3833	74	180	KU
2007 Feb 20	54152.1114	54152.3373	100	180	KU
2007 Feb 21	54153.2148	54153.2973	40	180	KU
2007 Feb 23	54155.0150	54155.1822	80	180	KU
2007 Feb 25	54157.3127	54157.3804	33	180	KU
2007 Feb 26	54158.2717	54158.3311	28	180	KU
2007 Feb 28	54160.1487	54160.2101	30	180	KU

* HJD - 2450000. [†] Number of frames.[‡] Exposure time in unit of second. [§]Observers' ID, see table 2.**Fig. 2.** The theta diagram of the PDM analysis. This provides the best estimated period of 0.069824 d.

Also shown in this figure is the dashed curve by fitting a quadratic equation between $E=0$ and 73. This is given by

$$O - C = -3.97(1.05) \times 10^{-3} - 8.57(6.66) \times 10^{-4}E + 1.95(0.90) \times 10^{-6}E^2. \quad (3)$$

From these quadratic equations, we obtained $P_{\text{dot}} = \dot{P}/P = 8.9(1.1) \times 10^{-5}$ from the former equation and $\dot{P}/P = 5.6(2.6) \times 10^{-5}$ from the latter equation, respectively. These results imply that the superhump period increased at $\dot{P}/P = 7(+3, -4) \times 10^{-5}$ in the middle of the plateau phase.

After $E=100$, the datapoints can be fitted with a linear equation. This indicates that the superhump period remains almost constant during this phase. A linear regression to the observed maximum timings between $E=100$ and 174 yields

$$HJD(max) = 2454139.2969(20) + 0.06954(1) \times E. \quad (4)$$

In conjunction with the $O - C$ diagram and the above equations, we found that a sudden period decrease oc-




Fig. 3. Daily averaged light curves of the 2007 February superoutburst. The numbers in this figure indicate HJD-2454100.




Fig. 4. Enlarged light curve obtained by KU on 2007 February 12. A characteristic feature of superhumps is visible. The absence of an eclipse means a low-to-mid inclination of NSV 4838.

Table 3. Timings of the superhump maxima.

E^*	HJD [†]
0	4139.2549
1	4139.3247
44	4142.3284
54	4143.0243
55	4143.0943
56	4143.1634
57	4143.2360
68	4144.0064
69	4144.0733
70	4144.1444
71	4144.2129
72	4144.2850
73	4144.3563
100	4146.2519
101	4146.3194
113	4147.1551
114	4147.2272
115	4147.2954
127	4148.1296
128	4148.1980
129	4148.2658
155	4150.0748
156	4150.1461
157	4150.2157
158	4150.2872
159	4150.3544
169	4151.0520
170	4151.1188
172	4151.2564
173	4151.3283

* Cycle count.

† HJD-2450000.

curred between $E=80$ and $E=100$.

3.2. 2005 June superoutburst

The light curves during the 2005 June outburst are shown in figure 6. From HJD 2453528 to HJD 2453532, the magnitude declined at a rate of $0.06(1) \text{ mag d}^{-1}$. Although this value is smaller compared to the average decline rate of the plateau stage (0.10 mag d^{-1} , Warner 1995), such a slow decline is sometimes observed at the late phase of the plateau stage (Baba et al. 2000; Patterson et al. 2000). Enlarged light curves are displayed in figure 7, in which hump-like features with an amplitude of 0.3 mag are visible. After subtracting the linear declining trends, we also performed the PDM for the data over the first 4 nights and found a weak signal at $0.0699(1)$ days. This value is in good accordance with that obtained during the 2007 February superoutburst. Judging from the above results and other information such as the AAVSO light curve generator⁴, we conclude that we observed NSV 4838 from the late stage of the superoutburst.

⁴ <http://www.aavso.org/data/lcg/>




Fig. 5. $O - C$ diagram of the superhump maxima. The horizontal and vertical axes denote the cycle count and $O - C$, respectively. The solid and dashed curve correspond to the equation (2) and (3), respectively. After $E=100$, the data points can be fitted with the equation (4), which indicates that the superhump period unchanged in this phase.




Fig. 6. Light curves of the 2005 June outburst obtained by JP. The horizontal and vertical axes indicate the HJD-2453500 and the relative magnitude, respectively.




Fig. 7. Light curves of the 2005 June superoutburst obtained by JP. The numbers in this figure indicate HJD-2453500. Although the data were noisy, superhumps are visible with an amplitude of 0.3 mag.

3.3. mass ratio

It is well known that the mass ratio of the system is related with the fractional superhump excess (Patterson 1998; Patterson et al. 2005; Knigge 2006). Here we use an empirical relation derived by Patterson (1998) as follows:

$$\epsilon = \frac{0.23q}{1 + 0.27q}, \quad (5)$$

where ϵ is the fractional period excess, and q is the mass ratio of the system, respectively. Using $P_{\text{sh}}=0.0698$ days and $P_{\text{orb}}=0.0678$ days (Thorstensen, unpublished, see also Ritter, Kolb 2003), we can estimate the mass ratio of NSV 4838 to be 0.13. This result is in good accordance with that using other relations such as the equation (8) of Patterson et al. (2005), equation (7) of Knigge (2006).

4. Discussion

4.1. long-term behavior of NSV 4838

The present photometric studies have shown the basic properties of NSV 4838. The mean superhump periods, the fractional superhump excess, and the duration of the plateau stage are quite a typical for the "textbook" SU UMa-type dwarf novae (Warner 1995). We also investigated the long-term behavior of NSV 4838 based mainly on the AAVSO light curve generator after the 2005 June superoutburst. The data include 229 negative observations and 104 positive observations from 2005 June to 2008 April. Although the short baseline of the archival data and its faintness even at the maximum brightness cannot give a convincing result as to whether some outbursts had been missed, we detected six outbursts in total. Figure 8 shows the long-term light curves of NSV 4838. If we missed no superoutbursts, an estimated supercycle of NSV 4838 is about 340 days, which is a typical value for SU UMa-type dwarf novae. On the other hand, the total number of the recorded normal outburst is too small for the estimated supercycle of NSV 4838 (Osaki 1995). One possibility may be that the faintness of its maximum magnitude of normal outburst prevented its detection. If we did not overlook any normal outbursts, the event ratio of the normal outburst to superoutburst is unusually small (see. e.g., table 1 of Osaki (1995)). Future continuous observations of this object are needed to clarify the activity of NSV 4838.

4.2. distance

According to an empirical relation of Harrison et al. (2004), the absolute magnitude of dwarf novae at the maximum of the *normal outburst* is given as a function of its orbital period,

$$M_V = 5.92 - 0.383 \times P_{\text{orb}}, \quad (6)$$

where P_{orb} is in the unit of hour. According to multicolor photometry, the color index shows $V - R_c \sim 0$ near the bright maximum (Uemura et al. 2008). Thus the maximum magnitude in V band may be almost the same value as that in R_c band. Since no filter is close to R_c , it is not




Fig. 8. Long-term light curves of NSV 4838 since the 2005 June superoutburst using the AAVSO archive. The filled circles and bottom triangles indicate positive and negative observations, respectively. For display purpose, we precluded negative observations brighter than 16.0 mag. We also designate superoutburst and normal outburst as “S” and “N”, respectively.

unreasonable that we regard the magnitude in non-filter as that in *V* band.

Extensive observations of SU UMa-type dwarf novae suggest that the maximum magnitude of the normal outburst is fainter than that of the superoutburst by about 0.5 mag (Warner 1995; Shafter et al. 2007). In conjunction with the above facts including figure 8, a maximum magnitude of normal outburst for NSV 4838 is expected to be around 15.5 mag in *V* band. Using this value, we roughly estimated 1.1 kpc as a distance to NSV 4838.

4.3. superhump period change

The general consensus concerning the superhump period change is that the majority of SU UMa-type dwarf novae show a decrease in their superhump periods during superoutburst (Uemura et al. 2005). Osaki (1985) suggested that this is possibly due to shrinkage of the disk radius, or consequence of mass depletion from the accretion disk. Recently, it has been recognized that some SU UMa-type dwarf novae show an increasing superhump period change during a plateau phase of superoutburst (Semeniuk et al. 1997; Nogami et al. 1997). So far, we have confirmed this increase for about 15 SU UMa-type dwarf novae, most of which have their superhump periods shorter than 0.065 days (Oizumi et al. 2007).

Based on the refined thermal-tidal instability model proposed by Osaki, Meyer (2003), the maximum radius during the superoutburst without a precursor is larger than that with a precursor. When the accretion disk reaches the tidal truncation radius (the maximum radius) at the onset of the outburst, the matter will be piled up at this radius, at which the stored matter works as a wall. The wall prevents the cooling wave from propagating inward because the continuous tidal dissipation at the tidal truncation radius urges matter to infall. In this case, no precursor is observed at the onset of the superoutburst. On the other hand, when the accretion disk does not reach the tidal truncation radius at the onset of the outburst, the cooling wave can propagate from the outermost an-

nulus, which is observed as a rapid decline like a normal outburst. If the eccentricity of the accretion disk grows to some extent, the heating wave can again propagate from the outermost annulus, because the tidal torque enhancement supplies the gas to the inner annulus. This process is observed as a superoutburst with a precursor.

Recently, Uemura et al. (2005) studied superhump period changes for the 2001 and 2004 superoutbursts of a short period SU UMa-type dwarf nova TV Crv. It was found that the 2001 superoutburst showed no evidence of a precursor and the superhump period increased during the plateau phase. On the other hand, the 2004 superoutburst was accompanied by a precursor and the superhump period kept almost constant during the plateau phase. In combination with this observations and the refined thermal-tidal instability model, Uemura et al. (2005) suggested that enough propagation of the eccentric mode outside the 3:1 resonance radius may be observed as an increase in the superhump period. Therefore, short period SU UMa-type dwarf novae, in which there exists a large annulus between the 3:1 resonance radius and tidal truncation radius, tend to show a positive period derivative. On the other hand, long period SU UMa-type dwarf novae and those with a precursor during superoutburst, in which the gap between the 3:1 resonance radius and the tidal truncation radius is small, tend to show negative or zero period derivatives.

Figure 9 illustrates the $P_{\text{dot}}-P_{\text{sh}}$ diagram of SU UMa-type novae, in which we added the results of NSV 4838. It should be noted that NSV 4838 showed the positive period derivatives despite the fact that the superhump period of the system is as long as 0.070 days. In figure 9, one can notice that a few systems of $P_{\text{sh}} > 0.07$ days also show the positive P_{dot} . Such systems include VW CrB ($P_{\text{sh}}=0.07287$ days, Nogami et al. 2004), TT Boo ($P_{\text{sh}}=0.07796$ days, Olech et al. 2004), and RZ Leo ($P_{\text{sh}}=0.07853$ days, Ishioka et al. 2001). Interestingly, these systems commonly show long-lasting (exceeding 2 weeks), large-amplitude (exceeding 6 mag) superoutbursts




Fig. 9. $P_{\text{dot}}-P_{\text{sh}}$ diagram of SU UMa-type dwarf novae. The location of NSV 4838 in this figure is designated as filled circles. The original data were taken from Oizumi et al. (2007) (See also Uemura et al. (2005) for a comprehensive table of P_{dot} values.).

and long supercycles. On the other hand, NSV 4838 exhibits relatively a small amplitude (~ 4 mag)⁵ and a moderate supercycle (~ 1 year). In addition, the negative P_{dot} was reported in the 1991 October superoutburst of EF Peg, which exhibited a long-lasting and large-amplitude superoutburst, as well as a long recurrence time (Howell et al. 1993; Kato 2002). Therefore, we suggest that these properties do not necessarily lead to the positive P_{dot} .

Although the exact mechanism causing superhump period changes still remains unknown, the present results imply that an additional physical parameter on the accretion disk should be required for explaining the observed diversity of the superhump period changes. In order to clarify the nature of the superhump period changes, we should collect further samples.

5. Summary

In this paper, we present CCD photometry of the dwarf nova NSV 4838 during the 2005 June and 2007 February superoutbursts, as well as the long-term behavior of the system. During the 2007 February superoutburst, the best estimated superhump period was 0.069824(83)d. We also examined superhump period changes and concluded that the superhump period increased with $\dot{P}/P = +7(+3, -4) \times 10^{-5}$. In the late phase of the superoutburst, the mean superhump period became almost constant at 0.06954(1) days. Based on the long-term light curves, we found four superoutbursts since 2005 June and estimated a supercycle of NSV 4838 to be about 340 days. On the other hand, we detected at most two normal outbursts. If we missed no normal outbursts, the event ratio of the normal outburst to the superoutburst is unusually small for this supercycle. Using the empirical relations, we derived

about 1.1 kpc as the distance to NSV 4838 and 0.13 as the mass ratio, respectively. We plotted the obtained values of NSV 4838 in the $P_{\text{dot}}-P_{\text{sh}}$ plane, finding that the positions in the plane deviate from the general trend. In order to understand the observed diversity of the superhump period changes, additional physics should be invoked. This should be clarified in future observations by collecting further information on superhump period changes.

We acknowledge with thanks the variable star observations from the AAVSO and VSNET International Database contributed by observers worldwide and used in this research. This work is supported by a Grant-in-Aid for the 21st Century COE “Center for Diversity and Universality in Physics” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT). This work is partly supported by a grant-in aid from the Ministry of Education, Culture, Sports, Science and Technology (No. 16340057, 17340055, 17740105, 18740153). Part of this work is supported by a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists (AI).

References

- Baba, H., Kato, T., Nogami, D., Hirata, R., Matsumoto, K., & Sadakane, K. 2000, PASJ, 52, 429
- Berger, J., & Fringant, A.-M. 1977, A&AS, 28, 123
- Berry, R., & Burnell, J. 2000, The handbook of astronomical image processing (The handbook of astronomical image processing, Richmond, VA: Willmann-Bell, 2000, xxviii, 624 p., 1 CD-ROM. ISGN 0943396670)
- Cannizzo, J. K., Shafter, A. W., & Wheeler, J. C. 1988, ApJ, 333, 227
- Cannon Smith, R. 2007, astro-ph/0701654
- Downes, R., Webbink, R. F., & Shara, M. M. 1997, PASP, 109, 345
- Downes, Ronald A., Webbink, Ronald F., Shara, Michael M., Ritter, Hans, Kolb, Ulrich, & Duerbeck, Hilmar W. 2001, PASP, 113, 764
- Fernie, J. D. 1989, PASP, 101, 225
- Gänsicke, B. T. 2005, in The Astrophysics of Cataclysmic Variables and Related Objects, ed. J.-M. Hameury, & J.-P. Lasota Vol. 330 of Astronomical Society of the Pacific Conference Series, 3
- Harrison, T. E., Johnson, J. J., McArthur, B. E., Benedict, G. F., Szkody, P., Howell, S. B., & Gelino, D. M. 2004, AJ, 127, 460
- Hellier, C. 2001, Cataclysmic Variable Stars: how and why they vary (Berlin: Springer-Verlag)
- Hirose, M., & Osaki, Y. 1990, PASJ, 42, 135
- Hoshi, R. 1979, Prog. Theor. Phys., 61, 1307
- Howell, S. B., Schmidt, R., DeYoung, J. A., Fried, R., Schmeer, P., & Gritz, L. 1993, PASP, 105, 579
- Ishioaka, R., et al. 2001, PASJ, 53, 905
- Kato, T. 2002, PASJ, 54, 87
- Knigge, C. 2006, MNRAS, 373, 484
- Meinunger, L. 1981, Inf. Bull. Variable Stars, 2001
- Murray, J. R. 1998, MNRAS, 297, 323
- Nogami, D., Kato, T., Baba, H., Matsumoto, K., Arimoto, J., Tanabe, K., & Ishikawa, K. 1997, ApJ, 490, 840

⁵ In dwarf novae during quiescence, V magnitude is close to g (e.g., Szkody et al. 2005). From this, V magnitude in quiescence may be around 18.8, which corresponds to 3.7 in figure 1.

- Nogami, D., Uemura, M., Ishioka, R., Kato, T., & Pietz, J. 2004, PASJ, 56, S155
- Oizumi, S., et al. 2007, PASJ, 59, 643
- Olech, A., Cook, L. M., Złoczewski, K., Mularczyk, K., Kędzierski, P., Udalski, A., & Wisniewski, M. 2004, Acta Astron., 54, 2330
- Osaki, Y. 1985, A&A, 144, 369
- Osaki, Y. 1989, PASJ, 41, 1005
- Osaki, Y. 1995, PASJ, 47, 47
- Osaki, Y. 1996, PASP, 108, 39
- Osaki, Y. 2005, Proceedings of the Japan Academy, Ser. B: Physical and Biological Sciences, Vol. 81, p. 291-305., 81, 291
- Osaki, Y., & Meyer, F. 2003, A&A, 401, 325
- Patterson, J. 1998, PASP, 110, 1132
- Patterson, J., et al. 2005, PASP, 117, 1204
- Patterson, J., et al. 2000, PASP, 112, 1584
- Ritter, H., & Kolb, U. 2003, A&A, 404, 301
- Semeniuk, I., Olech, A., Kwast, T., & Nalezyty, M. 1997, Acta Astron., 47, 201
- Shafter, A. W., Coelho, E. A., & Reed, J. K. 2007, PASP, 119, 388
- Smith, A. J., Haswell, C. A., Murray, J. R., Truss, M. R., & Foulkes, S. B. 2007, ArXiv e-prints, 704
- Stellingwerf, R. F. 1978, ApJ, 224, 953
- Szkody, P., et al. 2005, AJ, 129, 2386
- Uemura, M., et al. 2008, PASJ, 60, 227
- Uemura, M., et al. 2005, A&A, 432, 261
- Warner, B. 1995, Cataclysmic Variable Stars (Cambridge: Cambridge University Press)
- Whitehurst, R. 1988, MNRAS, 232, 35
- Yanagisawa, K., et al., 2006, SPIE, 6269, 118